

Head Lice


What are head lice?

Head lice are parasitic insects that can live in your hair, eyebrows, and eyelashes. They cannot reproduce without the warmth of the head, nor can they survive without the blood provided by the scalp. Head lice should not be confused with body lice or crab lice. They are found only on humans and not on dogs, cats, or other pets.

What are the signs and symptoms of head lice?

Infestation is defined as harboring any nits, nymphs, or adult head lice. Excessive scratching of the head is the most common symptom of head lice infestation. You may also experience a tickling feeling or sensation something is in the hair, irritability and sleeplessness, or sores on the head caused by scratching.

What do head lice look like?

Lice Life Cycle	Description	Illustration
Nits (Eggs)	Nits look like white or dark ovals and are most found on the back of the neck and around the ears attached firmly within 1/4 inch of the base of the hair shaft.	
Nymph	A nymph is an immature louse that hatches from the nit. A nymph looks like an adult louse but is smaller. Nymphs mature into adults about 9–12 days after hatching from the nit.	
Adult	Actual lice are seen infrequently as they move quickly through the hair and avoid light. Lice are about the size of a sesame seed, and while they can crawl, they cannot jump or fly.	

How are head lice spread?

Lice are spread through direct or indirect contact with infested objects or people. When combs or brushes are shared, lice can be transported from one head to another. This is also true for the sharing of hats and other personal items, or clothing that is used or work on the head. If lice or their eggs are living, they can be moved from person to person and cause infestation.

Can head lice cause any illnesses?

Head lice have not been associated with the transmission of any other diseases. However, bacterial infections can occur due to scratching of the scalp. If you believe you have an infection, follow up with your healthcare provider for the appropriate treatment.

How soon after exposure to lice would an infestation be noticed?

It might take two to three weeks to notice the itching associated with infestation, but nits can be seen before the itching begins.

Who should be treated for head lice?

Treatment for head lice is recommended for anyone diagnosed with an active infestation. All household members and other close contacts should be checked for active infestation and treated. Household members including bedmates of infested persons should be treated at the same time.

What is the treatment for head lice?

1. Consult with a nurse, pharmacist, or physician and purchase an over the counter or prescription treatment. Read enclosed product directions before initiating treatment. There are products called “pediculicides” which can be bought over the counter or with a prescription from a healthcare provider. Pediculicides contain chemicals that kill lice. These products should be used carefully after reading the directions thoroughly. It is important to remember that **NO PEDICULICIDE TREATMENT IS 100% EFFECTIVE**. Repeat treatments may be necessary.
2. It is important to remove all lice and nits by hand. You can do this using a lice comb or a comb with very fine teeth. Check all household members at the same time, so that everyone needing it can be treated at the same time.
3. Do not use a combination shampoo/conditioner, or conditioner before using lice medicine. Do not re-wash the hair for 1–2 days after the lice medicine is removed.
4. Have the infested person put on clean clothing after treatment.
5. If a few live lice are still found 8–12 hours after treatment, but are moving more slowly than before, do not retreat. The medicine may take longer to kill all the lice. Comb dead and any remaining live lice out of the hair using a fine-toothed nit comb.
6. If no dead lice are found after 8-12 hours and lice seem as active as before, the medicine may not be working. Talk to your health care provider for advice.
7. The home environment, such as bedding, clothing, etc., must also be treated at the same time.

Tips for Nit Removal

1. While hair is wet, separate into sections.
2. Comb through each section of hair and remove all nits with a lice comb.
3. Rinse hair and scalp with running water. Allow hair to air dry.
4. Once hair is dry, recheck entire head and remove any remaining nits.
5. Put on clean clothes.

Follow Up of Cases

1. Do daily head checks for the next 10 days, and remove any nits or lice found.
2. If no nits or lice are found on days 7-10, a second treatment with a lice-killing product is not necessary. If any lice or nits are still present on or after day 7, a second treatment should be performed following the steps outlined in the “Treatment” and “Nit Removal” sections. Continue with daily head checks for the next 10 days.
3. After the child is free of lice and nits for 10 days, continue head check as part of routine hygiene.

Treatment Precautions

1. Only use licensed and approved products for treatment of head lice. Home remedies such as mayonnaise, Vaseline, and tea tree oil are not consistently proven to be effective for the treatment of head lice. Tea tree oil is not to be used daily and is toxic to the liver in high doses. Pure tea tree oil can be irritating to the skin and can provoke an allergic reaction in some individuals.
2. Do not use any other products like gasoline, kerosene, petroleum jelly (Vaseline), or other products containing insecticides that are not labeled for use on humans to treat head lice.
3. The treatment times of over-the-counter lice shampoos and rinses must not be extended beyond the package insert recommendations or applied too frequently.

Treating the Environment

1. At the same time medicating shampooing and nit removal are performed, machine wash on the hot cycle (130° F or hotter) all bed linens and clothing that have been in contact with the infested person within the last three days. Also, wash the soft toys and stuffed animals that accompany the child to bed. Use a hot dryer setting for at least 20 minutes to dry clothes, linens, and soft/stuffed animals after washing.
2. All the child's brushes, combs, and hair accessories like barrettes, ponytail holders, and headbands must be treated as well. The following methods are suggested: a) Soak items in a mild bleach solution (one tablespoon of bleach per quart of cool water), rubbing alcohol, or Lysol for one hour, or b) Scrub items with soap and hot (130° F) water.
3. If there are items which cannot be wash, vacuumed or dry cleaned, items can be "bagged" and sealed in plastic bags for 2 weeks.
4. Non-washables can be vacuumed or dry-cleaned. Vacuum carpet, upholstered furniture, mattresses, box springs, and car seats.
5. The use of lice sprays, house "bombs", exterminator services, or treatment of household pets are not needed and could be harmful.

How can lice be prevented in school settings?

The Oklahoma State Department of Health does not advocate exclusion of children from school, but rather supports a cooperative effort between parents, the schools, local healthcare providers and public health personnel, local state human services agencies, and local childcare advocacy groups to prevent and control pediculosis. Children should not be excluded from school but should be sent home with educational materials for prompt treatment and nit removal. Head lice control policies may differ from one school district to another. Contact the school administrator or nurse to learn more about your child's school head lice policies.